

Systematic review of economic analyses of health care-associated infections

Patricia W. Stone, PhD, RN, Deborah Braccia, MPA, RN, and Elaine Larson, PhD, RN, FAAN, CIC
New York, New York

Background: Economic evidence is needed to assess the burden of health care-associated infections (HAIs) and cost-effectiveness of interventions aimed at reducing related morbidity and mortality. The objective of this study was conducted to assess the quality of economic evaluations related to HAI and synthesize the evidence.

Methods: A systematic review of research published between January 2001 and June 2004 was conducted. Quality of the publication was estimated using a Likert-type scale. All cost estimates were standardized into a common currency. Descriptive statistics and a logistic regression were conducted to identify predictors of high quality.

Results: 70 studies were audited. There was wide variation in these cost estimates. Publications estimating the cost attributable to an infection were almost 7 times more likely judged to be of higher quality than studies of the cost of interventions ($P < .05$). Papers in which the authors stated the perspective (hospital or societal) were twice as likely to be judged as being of high quality ($P < .05$).

Conclusion: There are more publications and growing interest in estimating the costs of HAI. However, the methods employed vary. We recommend (1) the use of guidelines for authors and editors on conducting an economic analysis, (2) development of more sophisticated mathematical models, and (3) training of infection control professionals in economic methods. (*Am J Infect Control* 2005;33:501-9.)

Health care-associated infections (HAI) are one of the most serious patient safety issues in health care today. In the United States, the incidence of HAI has been estimated to be approximately 2 million cases annually.^{1,2} More than 500,000 of these infections occur in intensive care units (ICUs), and most are associated with the presence of an invasive device such as a central venous catheter or ventilator. The rate of HAI per 1000 patient days increased 36% from 1975 to 1995.³ Furthermore, it has been estimated that there are approximately 90,000 deaths attributed to HAI annually, ranking it as the fifth leading cause of death in acute care hospitals.⁴ These trends suggest that many challenges still exist in the prevention and control of HAI in the health care setting.

The total annual hospital-related financial burden of HAI in the United States was estimated to exceed \$4.5 billion in 1992 (using the Consumer Price Inflator, this converts to 6.5 billion in 2004 dollars).¹ However,

we found no further national estimates of the burden, and this estimate is based on infection rates measured in the Study on the Efficacy of Nosocomial Infection Control (SENIC) study, which was conducted in the mid-1970s.⁵ Hence, these estimates are dated and likely to underestimate the current costs.

Increasingly, there are calls for the application of cost-effective technologies to decrease the burden of illness related to infectious diseases and specifically HAI.^{6,7} Therefore, it is critically important to understand both the economic burden of HAI as well as the evidence from cost-effectiveness research examining health-related technologies, services, and programs aimed at reducing HAI disease-related morbidity and mortality. To understand the economic burden of HAI, the resources and related costs of interest are the incremental costs that may be directly attributable to the infection and not the underlying admitting diagnosis. In a previous survey,⁸ we audited a decade of published economic evidence on the attributable costs of HAI and interventions aimed at reducing this burden. In that study, we found a wide variation in the cost estimates (eg, \$3500 to \$40,000 per survivor of bloodstream infection [BSI], in 2000 dollars).⁸ Differences in methods used to estimate costs, which has been a chronic problem found in economic evaluations, contributes to the wide range of costs reported.⁹⁻¹² In addition, results cited above include estimates of hospital costs only, not costs to the broader health care sector or to society. The objectives of our current study were to update that review and assess the quality of these analyses.

From the Columbia University School of Nursing, New York, NY.

Supported by an unrestricted grant from the Association of Professionals in Infection Control and Epidemiology Research Foundation through a grant from 3M Health Care.

Reprint requests: Patricia W. Stone, PhD, RN, Assistant Professor, Columbia University School of Nursing, 617 W. 168th St, New York, NY 10032. E-mail: ps2024@columbia.edu.

0196-6553/\$30.00

Copyright © 2005 by the Association for Professionals in Infection Control and Epidemiology, Inc.

doi:10.1016/j.ajic.2005.04.246

METHODS

A systematic review of existing published evidence was conducted. Articles were included if they were published between January 2001 and June 2004, had an abstract for review, contained an original cost estimate, and were written in English. To find the published analyses, searches were conducted in MEDLINE, EconoLit, and HealthSTAR using the medical subject headings (MeSH) or text keywords “nosocomial infections,” “infection control,” or “hospital acquired infections” cross referenced with “costs,” “cost analysis,” “economics,” or “cost-effectiveness analysis.” In addition, review articles were examined for published articles that met the inclusion criteria, and other published articles that were known to the authors were included. Methodologic articles and editorials were excluded.

Our original audit form containing 23 data elements was based on work from the Harvard Cost-Effectiveness Analysis Registry research team.¹³ We piloted and refined the form as needed. Data elements extracted included the type of study (simple cost analysis of HAI or economic evaluation of an intervention to prevent or reduce HAI), source of funding, country of study, type of HAI analyzed, and definition of HAI used (ie, Centers for Disease Control and Prevention [CDC] definition, other definition, or not stated) and whether the study included consideration of antibiotic resistance. Study design elements such as types of costs included, source of cost data, and source of effectiveness data or attributable costs (if applicable) were noted. If the study was an economic evaluation of an intervention, the type of economic evaluation (eg, simple cost analysis without comparison or cost-effectiveness analysis) was categorized. We also noted whether decision analytic models to estimate cost-effectiveness were used.

In accordance with methods recommended to audit systematically the economic evaluations,¹⁴ 2 trained readers audited each study and met for consensus discussions. If there was uncertainty or disagreement on any data element, a third expert reader was consulted. Similar to the Harvard Cost-Effectiveness Analysis registry audit, quality of the analysis was rated on a 7-point Likert scale, with the highest quality scored as 7 and the lowest quality assigned a 1. The quality score assigned to the analysis was based on the mean between the 2 readers. We then dichotomized this variable based on the distribution of data; studies with a mean score of 4 or less were considered low quality and greater than 4 were considered high quality.

Economic (ie, monetary) results were standardized into a common currency and year. When authors stated the year of currency in which their results were

reported, we used that year for calculations. If the year was not stated, 2 years prior to publication was used as an estimate of the probable year. All cost estimates were standardized into US currency with use of the Economic Research Federal Reserve Bank of St. Louis/FRED II (<http://research.stlouisfed.org/fred2/categories/15>) rates for January 1 of the estimated year of the study. In addition, all US dollars were standardized into 2002 values using the Bureau of Labor Statistics Consumer Price Index calculator (<http://data.bls.gov/cgi-bin/cpicalc.pl>).

Data were entered into an ACCESS Microsoft office database and analyzed using SPSS 11.5 (SPSS Inc, Chicago, IL). Descriptive statistics were computed. A logistic regression was conducted in which the quality of the analysis (high = a mean consensus score of greater than 4, and low = a mean consensus score of 4 or less) was the dependent variable and predictor variables included paper type (cost of infection or cost of intervention), country of study (United States or other), consideration of antibiotic resistance (yes or no), HAI definition (stated, not stated), perspective (stated, not stated), and source of funding (stated, not stated).

To summarize the results of studies, we grouped analyses by type of infection studied (urinary tract infections, ventilator associated pneumonias, BSI, surgical infections, and other). In this synthesis, results were only included if they were originally reported as cost per patient infection or if it was possible to convert the results to this standard outcome.

RESULTS

One hundred fifty-two manuscripts were obtained for review. Of these, 70 studies met our eligibility criteria. See [Appendix A](#) for full list of the eligible articles. [Table 1](#) summarizes publications in terms of country of study, funding source, description of HAI analyzed, and economic methods employed.

The majority of the analyses were conducted in the United States. Although the most commonly reported funding source for these studies was government or industry, many authors did not report whether there was or was not a funder. Surgical infections were most commonly analyzed, followed by BSI. Many studies examined organism-specific infections such as vancomycin-resistant enterococci or *Clostridium difficile*.

The most common analytic method was a simple cost analysis of infection, and the majority of the analyses were conducted from the perspective of the hospital. Costs related to outpatient or nonhealth care services were generally not included. Actual microcosting (eg, time and motion studies and/or costs of specific items) was the most frequent source of cost estimation.

Table 1. Cost analyses related to HAI January 2001 to June 2004

Paper characteristic	n	%	Economic methods	n	%
Country of study			Type of analysis		
United States	39	56	Cost analysis of infection	40	57
Europe	17	24	Cost analysis of intervention [†]	30	43
Australia/New Zealand	4	6			
Other	10	14	Perspective of analysis		
Source of funding*			Hospital	63	90
Government	14	20	Health care sector	6	9
Industry	13	19	Societal	1	1
Foundation	2	3	Costs included*		
In-kind	1	1	Intervention	25	36
Not stated	43	61	Hospitalization	60	86
HAI analyzed			Outpatient	3	4
Surgical	18	23.7	Antibiotics	9	13
Bloodstream	14	18.4	Nonhealth care	0	0
Pneumonia	6	7.9	Source of cost estimate*		
Urinary tract	2	2.6	Published data	14	20
Organism specific	25	32.9	Microcosting	50	71
Other	11	14.5	Estimated by authors	29	41
Antibiotic resistance considered			Claims	25	36
Yes	21	30	Other	3	4
Definition of HAI			Time horizon		
CDC	29	41	1 year or less	69	99
Other	31	44	Greater than 1 year	1	1
Not stated	10	14	Quality score		
			4 or less	35	50
			Greater than 4	35	50

N = 70.

*Categories are not mutually exclusive.

[†]Twenty of these studies had no comparator and therefore were not formal economic evaluations. Five of the studies were cost-effectiveness analyses, 3 were cost-consequence analyses, and 2 were cost-minimization analyses.

The results of the logistic regression are displayed in Table 2. The type of paper was a significant predictor of study quality, with publications estimating the cost attributable to an infection almost 7 times more likely judged to be higher quality than studies of the cost of interventions. Papers in which the authors stated the perspective (hospital or societal) were twice as likely to be judged high quality.

Forty-five simple cost analyses could be standardized into cost per patient infections, for a total of 47 separate results. Twenty-four of these studies examined a specific type of infection: surgical site, BSI, ventilator-associated pneumonia, and urinary tract infections. On average, BSIs were found to be the most expensive (Table 3); however, the standard deviations

Table 2. Predictors of high-quality studies*

	Odds ratio	95% CI
NI definition stated	7.6	0.1-4.5
Cost analysis of infection compared with cost analysis of intervention	6.5 [†]	1.8-23.4
Perspective stated	2.3 [†]	1.1-4.8
United States compared with other countries	3.0	0.8-11.1
Antibiotic resistance	1.7	0.4-6.3
Funding source stated	1.2	0.6-2.2

*High quality defined as a quality score greater than 4 out of a range from 1 to 7.

[†]P < .05.

of all infection types were quite large, indicating wide variations in the estimated costs per patient.

DISCUSSION

We have found substantial growth in published evidence estimating the cost of HAI and interventions aimed at decreasing the related morbidity and mortality. In our previous audit using the same inclusion criteria, there were only 55 eligible articles found over a 10-year period compared with the 70 publications we found over a two-and-a-half-year period.⁸ However, as in the previous audit, the majority of the articles were simple cost analyses conducted from the hospital rather than the societal perspective, and the methods used in these analyses were varied. As in our previous audit, BSIs were the most expensive with comparable wide variation in the estimated costs.

Over a two-and-a-half-year period, we found substantial growth in published evidence estimating the cost of HAI and interventions aimed at decreasing related morbidity and mortality. This growth may be related to the overall growth in publications, or it may reflect an increased interest to understand financial implications of HAI as well as increased pressure to demonstrate the cost-effectiveness of infection control departments. However, because of lack of standard methods, caution is needed when interpreting this growing body of evidence.

There has been a number of efforts to standardize the conduct and reporting of economic evaluations of health care technology.¹⁵⁻¹⁸ The Panel on Cost-Effectiveness Analysis in Health Care and Medicine convened by the US Public Health Service Department suggested a standard set of methodologic practices intended to improve the comparability of cost-effectiveness evidence, which is called a *Reference Case*.¹⁵⁻¹⁸ Some of the recommendations for an ideal reference case include adopting a societal perspective, reporting results in terms of dollars per quality adjusted life year (QALY) gained (called a cost-utility analysis), including downstream net costs (and savings), discounting future

Table 3. Attributable costs of HAI

Infection type	Attributable costs		Range		References
	Mean	SD	Minimum	Maximum	
Surgical site infection	25,546	39,875	1783	134,602	30-37
Bloodstream infection	36,441	37,078	1822	107,156	38-46
Ventilator-associated pneumonia	9969	2920	7904	12,034	47,48
Urinary tract infection	1006	503	650	1361	49,50

Only articles that could be summarized using the outcome cost per patient were used in this analysis.

costs and QALYs, and conducting a minimal standard set of sensitivity analyses (ie, analyses in which a parameter is varied to estimate the degree of influence it has on the results of the base analysis). In addition, the panel developed a checklist to be used for journal reports (Table 4).

Unfortunately, the analyses we audited were for the most part much less sophisticated and did not meet these recommended standards. Many of the studies only reported aggregate estimates of costs and could not be included in our synthesis (Table 3). Although the recommendations in Table 4 are more rigorous, at the very least, analysts should provide data and/or calculate costs per person to have a standard outcome across studies. Of note, economic analyses regarding infection control interventions should further consider externalities such as the impact on herd effects and herd immunity of communicable infectious disease interventions to prevent distortion of cost-effectiveness estimates.¹⁹

Other previous auditors of economic literature have also raised concerns about the quality of the published analyses.^{10-12,20} There is some evidence that the quality is improving,¹³ but improvement is clearly still needed in the cost studies related to HAI. Although the results of the individual analyses audited have potential implications for clinical decision making, guideline development, and resource allocation, the quality of these analyses should be improved if they are to have a wide impact on health policy.

There are several approaches that could be used to improve the quality of economic analyses. First, journal editors could adopt guidelines for reviewing economic studies. The *British Medical Journal* has developed such a guideline.^{21,22} The guidelines are grouped into 10 sections under 3 headings: study design, data collection, and analysis and interpretation of results. Under each section is a commentary outlining the reasons for the requirements and the main unresolved methodologic issues and explaining why firm guidelines cannot be given in some cases. The guidelines are similar to the recommendations in

Table 4. Checklist for Journal Report of Economic Evaluation

1. Framework
 - Background of the problem
 - General framing and design of the problem
 - Target population for the intervention
 - Other program descriptors
 - Description of comparator programs
 - Boundaries of the analysis
 - Time horizon
 - Statement of the perspective of the analysis
2. Data and methods
 - Description of event pathway
 - Identification of outcomes of interest in the analysis
 - Description of model used
 - Modeling assumptions
 - Diagram of event pathway/model
 - Software used
 - Complete information about the sources of effectiveness data, cost data, and preference weights
 - Methods for obtaining estimates of effectiveness, costs, and preferences
 - Critique of data quality
 - Statement of year costs
 - Statement of method used to adjust costs for inflation
 - Statement of type of currency
 - Sources and methods for obtaining expert judgment
 - Statement of discount rates
3. Results
 - Results of model validation
 - Base results (discounted and undiscounted): total costs and effectiveness, incremental costs and effectiveness, and incremental cost-effectiveness ratios
 - Results of sensitivity analyses
 - Other estimates of uncertainty, if available
 - Graphical representation of results
 - Aggregate cost and effectiveness information
 - Disaggregated results, as relevant
 - Secondary analyses using 5% discount rate
4. Discussion
 - Summary of reference case results
 - Summary of sensitivity analysis assumptions having important ethical implications
 - Limitations of the study
 - Relevance of the study results for specific policy questions or decisions
 - Results of related economic evaluations
5. Technical report in appendix or available on request

Adapted from Gold et al.¹⁶

Table 4. Editors could also develop a cadre of reviewers experienced in economic evaluation.

Second, we recommend continued development of sophisticated mathematical policy models. Mathematical models are used routinely to guide public policy decisions in many areas that affect human life and health. Environmental regulation and military planning and strategy are 2 areas in which models have gained stature as policy tools.²³⁻²⁵ Models are also routinely used in economic forecasting with implications for macroeconomic policy, in transportation

planning with implications for the location and operation of traffic controls and the design of roadways, and in many other areas.²⁶

The US government has a long history of using and developing models to guide public health policy. Models rather than direct evidence have been used to support vaccine recommendations by the CDC at least since the late 1960s. The CDC's report "An Ounce of Prevention" reviewed and endorsed a variety of model-based estimates of gains in quality-adjusted life expectancy and cost-effectiveness ratios.²⁷ The Institute of Medicine has twice issued recommendations to use models to recommend priorities for the development of new vaccines.^{28,29}

The development of mathematical models to assess cost-effectiveness of interventions requires specific expertise. In the summer of 2004, in an effort to help nurse researchers better understand these techniques, a 2-day workshop was held by the National Institute of Nursing Research. Expert faculty from the Agency for Healthcare Research and Quality (AHRQ), Johns Hopkins University, and Columbia University provided a series of lectures and breakout sessions to help the nurse researcher understand the mechanics of these types of models (<http://www.ijhn.jhmi.edu/CostEffectivenessAnalysis/overview.htm>). Developing such workshops for infection control professionals may increase the rigor of the economic evaluations found in this literature.

In summary, there is increased interest in the economic evidence regarding the attributable costs of HAI and the cost-effectiveness of interventions aimed at reducing the morbidity and mortality associated with HAI. This has resulted in a growing number of publications. Surgical and BSI are the most frequently studied infections and the most expensive. However, there is wide variation in the methods used, limiting the impact of these analyses. The quality of economic evaluations should be increased to inform better the decision makers and clinicians.

References

1. CDC. Public health focus: surveillance, prevention, and control of nosocomial infections. *MMWR Morb Mortal Wkly Rep* 1992;41:783-7.
2. Burke JP. Infection control: a problem for patient safety. *N Engl J Med* 2003;348:651-6.
3. CDC. Monitoring hospital-acquired infections to promote patient safety—United States, 1990-1999. *MMWR Morb Mortal Wkly Rep* 2000;49:149-53.
4. Weinstein RA. Nosocomial infection update. *Emerg Infect Dis* 1998;4:416-20.
5. Haley RW, Emori TG. The Employee health service and infection control in US hospitals, 1976-1977. II. Managing employee illness. *JAMA* 1981;246:962-6.
6. Fos PJ, Miller DL, Amy BW, Zuniga MA. Combining the benefits of decision science and financial analysis in public health management: a county-specific budgeting and planning model. *J Public Health Manag Pract* 2004;10:406-12.
7. Global IDEA Scientific Advisory Committee. Health and economic benefits of an accelerated program of research to combat global infectious diseases. *CMAJ* 2004;171:1203-8.
8. Stone PW, Larson E, Kavar LN. A systematic audit of economic evidence linking nosocomial infections and infection control interventions: 1990-2000. *Am J Infect Control* 2002;30:145-52.
9. Neumann PJ, Stone PW, Chapman RH, Sandberg EA, Bell CM. The quality of reporting in published cost-utility analyses, 1976-1997. *Ann Intern Med* 2000;132:964-72.
10. Gerard K. Cost-utility in practice: a policy maker's guide to the state of the art. *Health Policy* 1992;21:249-79.
11. Gerard K, Mooney G. QALY league tables: handle with care. *Health Econ* 1993;2:59-64.
12. Gerard K, Smoker I, Seymour J. Raising the quality of cost-utility analyses: lessons learnt and still to learn. *Health Policy* 1999;46:217-38.
13. Neumann PJ, Greenberg D, Olchanski N, Stone PW, Rosen A. Growth and quality of the cost-utility literature, 1976-2001. *Value Health* 2005;8:3-9.
14. Carande-Kulis VG, Maciosek MV, Briss PA, Teutsch SM, Zaza S, Truman BI, et al. Methods for systematic reviews of economic evaluations for the Guide to Community Preventive Services. Task Force on Community Preventive Services. *Am J Prev Med* 2000;18:75-91.
15. Russell LB, Gold MR, Siegel JE, Daniels N, Weinstein MC. The role of cost-effectiveness analysis in health and medicine. Panel on Cost-Effectiveness in Health and Medicine. *JAMA* 1996;276:1172-7.
16. Gold MR, Siegel JE, Russell LB, Weinstein MC. Cost-effectiveness in health and medicine. New York: Oxford University Press; 1996.
17. Siegel JE, Weinstein MC, Russell LB, Gold MR. Recommendations for reporting cost-effectiveness analyses. Panel on Cost-Effectiveness in Health and Medicine. *JAMA* 1996;276:1339-41.
18. Weinstein MC, Siegel JE, Gold MR, Kamlet MS, Russell LB. Recommendations of the Panel on Cost-effectiveness in Health and Medicine. *JAMA* 1996;276:1253-8.
19. Brisson M, Edmunds WJ. Economic evaluation of vaccination programs: the impact of herd-immunity. *Med Decis Making* 2003;23:76-82.
20. Neumann PJ. Why don't Americans use cost-effectiveness analysis? *Am J Manag Care* 2004;10:308-12.
21. Drummond MF, Jefferson TO. Guidelines for authors and peer reviewers of economic submissions to the BMJ. The BMJ Economic Evaluation Working Party. *BMJ* 1996;313:275-83.
22. Smith R. New BMJ policy on economic evaluations. *BMJ* 2002;325:1124.
23. Weinstein MC, Toy EL, Sandberg EA, Neumann PJ, Evans JS, Kuntz KM, et al. Modeling for health care and other policy decisions: uses, roles, and validity. *Value Health* 2001;4:348-61.
24. Jones R. Use of modeling in developing label restrictions for agricultural chemicals. *Weed Technol* 1992;6:683-7.
25. Zubkoff P. The use of runoff and surface water transport and fate models in the pesticide regulation process. *Weed Technol* 1992;6:743-8.
26. National Research Council. Improving information for social policy decisions: the uses of microsimulation modeling. Vol. I. Review and recommendations. Washington, DC: National Academy Press; 1991.
27. Messonnier ML, Corso PS, Teutsch SM, Haddix AC, Harris JR. An ounce of prevention ... what are the returns? Second edition, 1999. *Am J Prev Med* 1999;16:248-63.

28. Institute of Medicine. New vaccines development: establishing priorities. Vol. I. Diseases of importance in the United States. Washington, DC: National Academy Press; 1985.
29. Institute of Medicine. Vaccines for the 21st Century: a tool for decision making. Washington, DC: National Academy Press; 1999.
30. Babcock HM, Carroll C, Matava M, L'Ecuyer P, Fraser V. Surgical site infections after arthroscopy: outbreak investigation and case control study. *Arthroscopy* 2003;19:172-81.
31. de la Torre SH, Mandel L, Goff BA. Evaluation of postoperative fever: usefulness and cost-effectiveness of routine workup. *Am J Obstet Gynecol* 2003;188:1642-7.
32. Dietrich ES, Schubert B, Ebner W, Daschner F. Cost efficacy of tazobactam/piperacillin versus imipenem/cilastatin in the treatment of intra-abdominal infection. *Pharmacoeconomics* 2001;19:79-94.
33. Engemann JJ, Carmeli Y, Cosgrove SE, Fowler VG, Bronstein MZ, Trivette SL, et al. Adverse clinical and economic outcomes attributable to methicillin resistance among patients with *Staphylococcus aureus* surgical site infection. *Clin Infect Dis* 2003;36:592-8.
34. Hollenbeak CS, Murphy D, Dunagan WC, Fraser VJ. Nonrandom selection and the attributable cost of surgical-site infections. *Infect Control Hosp Epidemiol* 2002;23:177-82.
35. Perencevich EN, Sands KE, Cosgrove SE, Guadagnoli E, Meara E, Platt R. Health and economic impact of surgical site infections diagnosed after hospital discharge. *Emerg Infect Dis* 2003;9:196-203.
36. Zhan C, Miller MR. Excess length of stay, charges, and mortality attributable to medical injuries during hospitalization. *JAMA* 2003;290:1868-74.
37. Whitehouse JD, Friedman ND, Kirkland KB, Richardson WJ, Sexton DJ. The impact of surgical-site infections following orthopedic surgery at a community hospital and a university hospital: adverse quality of life, excess length of stay, and extra cost. *Infect Control Hosp Epidemiol* 2002;23:183-9.
38. Bates DW, Yu DT, Black E, Sands KE, Schwartz JS, Hibberd PL, et al. Resource utilization among patients with sepsis syndrome. *Infect Control Hosp Epidemiol* 2003;24:62-70.
39. Diekema DJ, Beekmann SE, Chapin KC, Morel KA, Munson E, Doern GV. Epidemiology and outcome of nosocomial and community-onset bloodstream infection. *J Clin Microbiol* 2003;41:3655-60.
40. Dimick JB, Pelz RK, Consonni R, Swoboda SM, Hendrix CW, Lipsett PA. Increased resource use associated with catheter-related bloodstream infection in the surgical intensive care unit. *Arch Surg* 2001;136:229-34.
41. Johnson DW, MacGinley R, Kay TD, Hawley CM, Campbell SB, Isbel NM, et al. A randomized controlled trial of topical exit site mupirocin application in patients with tunnelled, cuffed haemodialysis catheters. *Nephrol Dial Transplant* 2002;17:1802-7.
42. Liu JV, Su YK, Liu CF, Chen JB. Nosocomial blood-stream infection in patients with end-stage renal disease: excess length of hospital stay, extra cost and attributable mortality. *J Hosp Infect* 2002;50:224-7.
43. Orsi GB, Di Stefano L, Noah N. Hospital-acquired, laboratory-confirmed bloodstream infection: increased hospital stay and direct costs. *Infect Control Hosp Epidemiol* 2002;23:190-7.
44. Rosenthal VD, Guzman S, Migone O, Crnich CJ. The attributable cost, length of hospital stay, and mortality of central line-associated bloodstream infection in intensive care departments in Argentina: a prospective, matched analysis. *Am J Infect Control* 2003;31:475-80.
45. Song X, Srinivasan A, Plaut D, Perl TM. Effect of nosocomial vancomycin-resistant enterococcal bacteremia on mortality, length of stay, and costs. *Infect Control Hosp Epidemiol* 2003;24:251-6.
46. Wisplinghoff H, Cornely OA, Moser S, Bethe U, Stutzer H, Salzberger B, et al. Outcomes of nosocomial bloodstream infections in adult neutropenic patients: a prospective cohort and matched case-control study. *Infect Control Hosp Epidemiol* 2003;24:905-11.
47. Dietrich ES, Demmler M, Schulgen G, Fekec K, Mast O, Pelz K, et al. Nosocomial pneumonia: a cost-of-illness analysis. *Infection* 2002;30:61-7.
48. Warren DK, Shukla SJ, Olsen MA, Kollef MH, Hollenbeak CS, Cox MJ, et al. Outcome and attributable cost of ventilator-associated pneumonia among intensive care unit patients in a suburban medical center. *Crit Care Med* 2003;31:1312-7.
49. Lai KK, Baker SP, Fontecchio SA. Impact of a program of intensive surveillance and interventions targeting ventilated patients in the reduction of ventilator-associated pneumonia and its cost-effectiveness. *Infect Control Hosp Epidemiol* 2003;24:859-63.
50. Tambyah PA, Knasinski V, Maki DG. The direct costs of nosocomial catheter-associated urinary tract infection in the era of managed care. *Infect Control Hosp Epidemiol* 2002;23:27-31.

Appendix A. Studies audited

1. Al-Ghamdi S, Gedebeu M, Bilal NE. Nosocomial infections and misuse of antibiotics in a provincial community hospital, Saudi Arabia. *J Hosp Infect* 2002;50:115-21.
2. Babcock HM, Carroll C, Matava M, L'Ecuyer P, Fraser V. Surgical site infections after arthroscopy: outbreak investigation and case control study. *Arthroscopy* 2003;19:172-81.
3. Bates DW, Yu DT, Black E, Sands KE, Schwartz JS, Hibberd PL, et al. Resource utilization among patients with sepsis syndrome. *Infect Control Hosp Epidemiol* 2003;24:62-70.
4. Bennett CL, Hynes D, Godwin J, Stinson TJ, Golub RM, Appelbaum FR. Economic analysis of granulocyte colony stimulating factor as adjunct therapy for older patients with acute myelogenous leukemia (AML): estimates from a Southwest Oncology Group clinical trial. *Cancer Invest* 2001;19:603-10.
5. Berild D, Ringertz SH, Lelek M, Fosse B. Antibiotic guidelines lead to reductions in the use and cost of antibiotics in a university hospital. *Scand J Infect Dis* 2001;33:63-7.
6. Brun-Buisson C, Roudot-Thoraval F, Girou E, Grenier-Sennelier C, Durand-Zaleski I. The costs of septic syndromes in the intensive care unit and influence of hospital-acquired sepsis. *Intensive Care Med* 2003;29:1464-71.
7. Capitano B, Leshem OA, Nightingale CH, Nicolau DP. Cost effect of managing methicillin-resistant *Staphylococcus aureus* in a long-term care facility. *J Am Geriatr Soc* 2003;51:10-6.
8. Carmeli Y, Eliopoulos G, Mozaffari E, Samore M. Health and economic outcomes of vancomycin-resistant enterococci. *Arch Intern Med* 2002;162:2223-8.
9. Church DL, Davies HD, Mitton C, et al. Clinical and economic evaluation of rapid influenza A virus testing in nursing homes in Calgary, Canada. *Clin Infect Dis* 2002;34:790-5.

10. de la Torre SH, Mandel L, Goff BA. Evaluation of post-operative fever: usefulness and cost-effectiveness of routine workup. *Am J Obstet Gynecol* 2003; 188:1642-7.
11. Diekema DJ, Beekmann SE, Chapin KC, Morel KA, Munson E, Doern GV. Epidemiology and outcome of nosocomial and community-onset bloodstream infection. *J Clin Microbiol* 2003;41:3655-60.
12. Dietrich ES, Demmler M, Schulgen G, et al. Nosocomial pneumonia: a cost-of-illness analysis. *Infection* 2002;30:61-7.
13. Dietrich ES, Schubert B, Ebner W, Daschner F. Cost efficacy of tazobactam/piperacillin versus imipenem/cilastatin in the treatment of intra-abdominal infection. *Pharmacoeconomics* 2001; 19:79-94.
14. Dimick JB, Pelz RK, Consunji R, Swoboda SM, Hendrix CW, Lipsett PA. Increased resource use associated with catheter-related bloodstream infection in the surgical intensive care unit. *Arch Surg* 2001;136:229-34.
15. Dimick JB, Swoboda SM, Pronovost PJ, Lipsett PA. Effect of nurse-to-patient ratio in the intensive care unit on pulmonary complications and resource use after hepatectomy. *Am J Crit Care* 2001;10:376-82.
16. Dominguez TE, Chalom R, Costarino AT Jr. The impact of adverse patient occurrences on hospital costs in the pediatric intensive care unit. *Crit Care Med* 2001;29:169-74.
17. Drinka PJ, Krause P, Nest L, Dissing M, Gravenstein S, Goodman BM. The effect of culture-positive influenza type A on resource use and adverse events in nursing home residents. *J Am Geriatr Soc* 2002;50:1416-20.
18. Engemann JJ, Carmeli Y, Cosgrove SE, et al. Adverse clinical and economic outcomes attributable to methicillin resistance among patients with *Staphylococcus aureus* surgical site infection. *Clin Infect Dis* 2003;36:592-8.
19. Fruhwirth M, Berger K, Ehlken B, Moll-Schuler I, Brosl S, Mutz I. Economic impact of community- and nosocomially acquired rotavirus gastroenteritis in Austria. *Pediatr Infect Dis J* 2001;20:184-8.
20. Geissler A, Gerbeaux P, Granier I, Blanc P, Facon K, Durand-Gasselin J. Rational use of antibiotics in the intensive care unit: impact on microbial resistance and costs. *Intensive Care Med* 2003;29:49-54.
21. Graves N, Nicholls TM, Morris AJ. Modeling the costs of hospital-acquired infections in New Zealand. *Infect Control Hosp Epidemiol* 2003; 24:214-23.
22. Harrington M, Butler K, Cafferkey M. Rotavirus infection in hospitalised children: incidence and impact on health care resources. *Ir J Med Sci* 2003;172:33-6.
23. Herr CE, Heckrodt TH, Hofmann FA, Schnettler R, Eikmann TF. Additional costs for preventing the spread of methicillin-resistant *Staphylococcus aureus* and a strategy for reducing these costs on a surgical ward. *Infect Control Hosp Epidemiol* 2003;24:673-8.
24. Hollenbeak CS, Alfrey EJ, Sheridan K, Burger TL, Dillon PW. Surgical site infections following pediatric liver transplantation: risks and costs. *Transpl Infect Dis* 2003;5:72-8.
25. Hollenbeak CS, Murphy D, Dunagan WC, Fraser VJ. Nonrandom selection and the attributable cost of surgical-site infections. *Infect Control Hosp Epidemiol* 2002;23:177-82.
26. Jenney AW, Harrington GA, Russo PL, Spelman DW. Cost of surgical site infections following coronary artery bypass surgery. *ANZ J Surg* 2001;71:662-4.
27. Johnson DW, MacGinley R, Kay TD, et al. A randomized controlled trial of topical exit site mupirocin application in patients with tunnelled, cuffed haemodialysis catheters. *Nephrol Dial Transplant* 2002;17:1802-7.
28. Karchmer TB, Durbin LJ, Simonton BM, Farr BM. Cost-effectiveness of active surveillance cultures and contact/droplet precautions for control of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect* 2002;51:126-32.
29. Kayihura V, Osman NB, Bugalho A, Bergstrom S. Choice of antibiotics for infection prophylaxis in emergency cesarean sections in low-income countries: a cost-benefit study in Mozambique. *Acta Obstet Gynecol Scand* 2003;82:636-41.
30. Kim T, Oh PI, Simor AE. The economic impact of methicillin-resistant *Staphylococcus aureus* in Canadian hospitals. *Infect Control Hosp Epidemiol* 2001;22:99-104.
31. Kyne L, Hamel MB, Polavaram R, Kelly CP. Health care costs and mortality associated with nosocomial diarrhea due to *Clostridium difficile*. *Clin Infect Dis* 2002;34:346-53.
32. Lai KK, Baker SP, Fontecchio SA. Impact of a program of intensive surveillance and interventions targeting ventilated patients in the reduction of ventilator-associated pneumonia and its cost-effectiveness. *Infect Control Hosp Epidemiol* 2003;24:859-63.
33. Lai KK, Fontecchio SA. Use of silver-hydrogel urinary catheters on the incidence of catheter-associated urinary tract infections in hospitalized patients. *Am J Infect Control* 2002;30:221-5.
34. Lemmen SW, Becker G, Frank U, Daschner FD. Influence of an infectious disease consulting service on quality and costs of antibiotic prescriptions in a university hospital. *Scand J Infect Dis* 2001; 33:219-21.

35. Liu JW, Su YK, Liu CF, Chen JB. Nosocomial bloodstream infection in patients with end-stage renal disease: excess length of hospital stay, extra cost and attributable mortality. *J Hosp Infect* 2002; 50:224-7.
36. Lopez H, Li JZ, Balan DA, et al. Hospital resource use and cost of treatment with linezolid versus teicoplanin for treatment of serious gram-positive bacterial infections among hospitalized patients from South America and Mexico: results from a multicenter trial. *Clin Ther* 2003;25:1846-71.
37. McNabb JJ, Nightingale CH, Quintiliani R, Nicolau DP. Cost-effectiveness of ceftazidime by continuous infusion versus intermittent infusion for nosocomial pneumonia. *Pharmacotherapy* 2001;21: 549-55.
38. Meyer NL, Hosier KV, Scott K, Lipscomb GH. Cefazolin versus cefazolin plus metronidazole for antibiotic prophylaxis at cesarean section. *South Med J* 2003;96:992-5.
39. Mokrzycki MH, Singhal A. Cost-effectiveness of three strategies of managing tunnelled, cuffed haemodialysis catheters in clinically mild or asymptomatic bacteraemias. *Nephrol Dial Transplant* 2002;17:2196-203.
40. Montecalvo MA, Jarvis WR, Uman J, et al. Costs and savings associated with infection control measures that reduced transmission of vancomycin-resistant enterococci in an endemic setting. *Infect Control Hosp Epidemiol* 2001;22:437-42.
41. Muto CA, Giannetta ET, Durbin LJ, Simonton BM, Farr BM. Cost-effectiveness of perirectal surveillance cultures for controlling vancomycin-resistant *Enterococcus*. *Infect Control Hosp Epidemiol* 2002; 23:429-35.
42. Nardi G, Di Silvestre AD, De Monte A, et al. Reduction in gram-positive pneumonia and antibiotic consumption following the use of a SDD protocol including nasal and oral mupirocin. *Eur J Emerg Med* 2001;8:203-14.
43. Oncul O, Yuksel F, Altunay H, Acikel C, Celikoz B, Cavuslu S. The evaluation of nosocomial infection during 1-year-period in the burn unit of a training hospital in Istanbul, Turkey. *Burns* 2002;28: 738-44.
44. Onen A, Cigdem MK, Geyik MF, et al. Epidemiology and control of nosocomial infections in paediatric surgery. *J Hosp Infect* 2002;52:166-70.
45. Orsi GB, Di Stefano L, Noah N. Hospital-acquired, laboratory-confirmed bloodstream infection: increased hospital stay and direct costs. *Infect Control Hosp Epidemiol* 2002;23:190-7.
46. Parodi S, Rhew DC, Goetz MB. Early switch and early discharge opportunities in intravenous vancomycin treatment of suspected methicillin-resistant staphylococcal species infections. *J Manag Care Pharm* 2003;9:317-26.
47. Pelz RK, Lipsett PA, Swoboda SM, et al. Vancomycin-sensitive and vancomycin-resistant enterococcal infections in the ICU: attributable costs and outcomes. *Intensive Care Med* 2002;28:692-7.
48. Perencevich EN, Sands KE, Cosgrove SE, Guadagnoli E, Meara E, Platt R. Health and economic impact of surgical site infections diagnosed after hospital discharge. *Emerg Infect Dis* 2003;9: 196-203.
49. Piednoir E, Bessaci K, Bureau-Chalot F, et al. Economic impact of healthcare-associated rotavirus infection in a paediatric hospital. *J Hosp Infect* 2003;55:190-5.
50. Piednoir E, Bureau-Chalot F, Merle C, Gotzamanis A, Wuibout J, Bajolet O. Direct costs associated with a nosocomial outbreak of adenoviral conjunctivitis infection in a long-term care institution. *Am J Infect Control* 2002;30:407-10.
51. Plowman R, Graves N, Griffin MA, et al. The rate and cost of hospital-acquired infections occurring in patients admitted to selected specialties of a district general hospital in England and the national burden imposed. *J Hosp Infect* 2001;47:198-209.
52. Puzniak LA, Gillespie KN, Leet T, Kollef M, Mundy LM. A cost-benefit analysis of gown use in controlling vancomycin-resistant *Enterococcus* transmission: is it worth the price? *Infect Control Hosp Epidemiol* 2004;25:418-24.
53. Roberts RR, Scott RD II, Cordell R, et al. The use of economic modeling to determine the hospital costs associated with nosocomial infections. *Clin Infect Dis* 2003;36:1424-32.
54. Rosenthal VD, Guzman S, Migone O, Crnich CJ. The attributable cost, length of hospital stay, and mortality of central line-associated bloodstream infection in intensive care departments in Argentina: a prospective, matched analysis. *Am J Infect Control* 2003;31:475-80.
55. Saizy-Callaert S, Causse R, Furhman C, Le Paih MF, Thebault A, Chouaid C. Impact of a multidisciplinary approach to the control of antibiotic prescription in a general hospital. *J Hosp Infect* 2003;53: 177-82.
56. Song X, Srinivasan A, Plaut D, Perl TM. Effect of nosocomial vancomycin-resistant enterococcal bacteremia on mortality, length of stay, and costs. *Infect Control Hosp Epidemiol* 2003;24: 251-6.
57. Spelman D, Harrington G, Russo P, Wesselingh S. Clinical, microbiological, and economic benefit of a change in antibiotic prophylaxis for cardiac surgery. *Infect Control Hosp Epidemiol* 2002;23: 402-4.

58. Stone PW, Gupta A, Loughrey M, et al. Attributable costs and length of stay of an extended-spectrum β -lactamase-producing *Klebsiella pneumoniae* outbreak in a neonatal intensive care unit. *Infect Control Hosp Epidemiol* 2003;24:601-6.
59. Tambyah PA, Knasinski V, Maki DG. The direct costs of nosocomial catheter-associated urinary tract infection in the era of managed care. *Infect Control Hosp Epidemiol* 2002;23:27-31.
60. Volkova N, Klapper E, Pepkowitz SH, Denton T, Gillaspie G, Goldfinger D. A case-control study of the impact of WBC reduction on the cost of hospital care for patients undergoing coronary artery bypass graft surgery. *Transfusion* 2002;42:1123-6.
61. Vriens M, Blok H, Fluit A, Troelstra A, Van Der Werken C, Verhoef J. Costs associated with a strict policy to eradicate methicillin-resistant *Staphylococcus aureus* in a Dutch University Medical Center: a 10-year survey. *Eur J Clin Microbiol Infect Dis* 2002;21:782-6.
62. Wahl WL, Franklin GA, Brandt MM, et al. Does bronchoalveolar lavage enhance our ability to treat ventilator-associated pneumonia in a trauma-burn intensive care unit? *J Trauma* 2003;54:633-9.
63. Warren DK, Shukla SJ, Olsen MA, et al. Outcome and attributable cost of ventilator-associated pneumonia among intensive care unit patients in a suburban medical center. *Crit Care Med* 2003;31:1312-7.
64. Webb M, Riley LW, Roberts RB. Cost of hospitalization for and risk factors associated with vancomycin-resistant *Enterococcus faecium* infection and colonization. *Clin Infect Dis* 2001;33:445-52.
65. Whitehouse JD, Friedman ND, Kirkland KB, Richardson WJ, Sexton DJ. The impact of surgical-site infections following orthopedic surgery at a community hospital and a university hospital: adverse quality of life, excess length of stay, and extra cost. *Infect Control Hosp Epidemiol* 2002;23:183-9.
66. Wisplinghoff H, Cornely OA, Moser S, et al. Outcomes of nosocomial bloodstream infections in adult neutropenic patients: a prospective cohort and matched case-control study. *Infect Control Hosp Epidemiol* 2003;24:905-11.
67. Zack JE, Garrison T, Trovillion E, et al. Effect of an education program aimed at reducing the occurrence of ventilator-associated pneumonia. *Crit Care Med* 2002;30:2407-12.
68. Zanetti G, Goldie SJ, Platt R. Clinical consequences and cost of limiting use of vancomycin for perioperative prophylaxis: example of coronary artery bypass surgery. *Emerg Infect Dis* 2001;7:820-7.
69. Zhan C, Miller MR. Excess length of stay, charges, and mortality attributable to medical injuries during hospitalization. *JAMA* 2003;290:1868-74.
70. Zhu XL, Wong WK, Yeung WM, et al. A randomized, double-blind comparison of ampicillin/sulbactam and ceftriaxone in the prevention of surgical-site infections after neurosurgery. *Clin Ther* 2001;23:1281-91.

Availability of Journal back issues

As a service to our subscribers, copies of back issues of *AJIC: American Journal of Infection Control* for the preceding 5 years are maintained and are available for purchase from Mosby until inventory is depleted. Please write to Mosby Subscription Customer Service, 6277 Sea Harbor Dr, Orlando, FL 32887, or call 800-654-2452 or 407-345-4000 for information on availability and prices of particular issues.